

HEALTHCARE PROVIDER INFORMATION

Name: _____
Address: _____
City: _____
State: _____ ZIP: _____
Phone: _____ Fax: _____

SUFLAVE[®]
(polyethylene glycol 3350,
sodium sulfate, potassium chloride,
magnesium sulfate, and sodium
chloride for oral solution)
178.7 g/7.3 g/1.12 g/0.9 g/0.5 g



SUFLAVE Dosing Regimen

SUFLAVE is a split-dose (2-day) regimen. A total of 2 bottles are required for complete preparation for colonoscopy. You will take 2 bottles of liquid in two separate doses. One dose of SUFLAVE is equal to one bottle plus one flavor enhancing packet.

For representational purposes only.

Day 1, Dose 1: Early in the Evening Before Your Colonoscopy

Start Dose 1 at _____ PM.

STEP 1 Open 1 flavor enhancing packet and pour the contents into 1 bottle.

STEP 2 Fill the provided bottle with lukewarm water up to the fill line. After capping the bottle, gently shake the bottle until all powder has mixed well (dissolved). For best taste, refrigerate the solution for an hour before drinking. Do not freeze. Use within 24 hours.

STEP 3 Drink 8 ounces of solution every 15 minutes until the bottle is empty.

STEP 4 Drink an additional 16 ounces of water during the evening.

IMPORTANT: If nausea, bloating, or abdominal cramping occurs, pause or slow the rate of drinking the solution and additional water until symptoms diminish.



For representational purposes only.


Day 2, Dose 2: The Morning of the Colonoscopy (5 to 8 hours prior to the colonoscopy and no sooner than 4 hours from starting Dose 1)

Start Dose 2 at _____ AM.

STEP 1 Repeat Step 1 to Step 3 from Day 1, Dose 1.

STEP 2 Drink an additional 16 ounces of water during the morning.

IMPORTANT: Continue to consume only clear liquids until after the colonoscopy. Stop drinking liquids at least 2 hours prior to colonoscopy.



For representational purposes only.



FINISH

Please see the other side for additional instructions.

